

New Appearance of Cosmic Coincidence Problem

Arthur D. Chernin

Sternberg Astronomical Institute, Moscow University, Moscow, 119899, Russia,

Astronomy Division, FIN-90014 University of Oulu, Finland,

and Tuorla Observatory, University of Turku, Piikkiö, FIN-21500, Finland

Abstract

The integrals of the Friedmann cosmology equations are identified as constant physical characteristics for both vacuum and non-vacuum cosmic energies. The integrals are found to be numerically coincident. A model shows that the coincidence can naturally originate in the electroweak scale physics at TeV temperatures. The coincidence seems to reflect a new type of symmetry that relates vacuum and non-vacuum energies.

PACS numbers: 98.80.Cq

Introduction. — Recent observations suggest that the cosmological expansion is accelerated [1,2]; this is found in the brightness-redshift distribution of high-redshift type-Ia supernovae [1] and confirmed by the bulk of all observational evidence that come from the cosmic age, large scale structure, and most convincingly from the combination of the cosmic microwave background (CMB) anisotropy and cluster dynamics [2,3]. It is assumed that cosmic acceleration is produced by cosmological vacuum, which has constant energy density $\rho_V \simeq 2 \times 10^{-123} M_{Pl}^4$, where $M_{Pl} \simeq 1 \times 10^{19}$ GeV is the Planck

energy scale.

In the Friedmann models, both energy density and pressure control cosmological expansion, so that the sum $\rho+3p$ presents the effective gravitating density in an isotropic universe. When the sum is negative, gravity becomes anti-gravity and accelerates the expansion. It is the case for vacuum, since vacuum has positive density ρ_V and negative pressure $p_V = -\rho_V$. Vacuum accelerates the expansion, if it dominates over ‘ordinary’ cosmic energies at the present era. The vacuum density measured in the units of the critical density is $\Omega_V = 0.7 \pm 0.1$, while the dark energy density is $\Omega_D = 0.3 \pm 0.1$, the baryon density is $\Omega_B = 0.02 \pm 0.01$, and the ultra-relativistic energy (CMB photons, gravitons, etc.) is $\Omega_R = 0.6\alpha \times 10^{-4}$ (where $1 < \alpha < 30 - 100$ is a dimensionless factor that accounts for possible non-CMB contributions to the energy) [2,3].

The density of vacuum is indeed larger than the total density of the non-vacuum energies. But surprisingly enough, the difference of the densities is not too big, and all the four are rather close to each other, especially the densities of vacuum and dark energy. It may mean that the vacuum started to dominate over the other energies not long ago, and so the present era is the era of transition from dark energy domination to vacuum domination. But why are all the four major densities coincident now? What is the physical nature of this phenomenon? This is the ‘cosmic coincidence problem’ that poses a severe challenge to current cosmological concepts (see [2,3] and references therein).

One possible approach to the problem would be to assume that the acceleration of the expansion is produced by non-vacuum energy which has negative pressure and negative effective gravitating density [4]. This energy form is called quintessence; it can naturally be realized in some scalar field models in which the field depends on time, and so the density of quintessence is diluted with the expansion. If so, the densities involved in the observed cosmic coincidence are all diluted, which makes them seemingly more similar to each other. Quintessence density might temporarily or even always be comparable to the rest of the energy densities [5]. Although such an idea may make the closeness of all cosmic densities natural, it does not explain the coincidence that the quintessence field becomes settled with a finite energy density comparable to the matter energy density

just now (see, for instance, [6]).

Another perspective on the cosmic coincidence problem is suggested by a model of Ref. [6] which proposes that the constant vacuum density may be determined in terms of two fundamental energy scales: the ‘reduced’ Planck scale $m_{Pl} \sim 10^{18}$ GeV and the electroweak scale $M_{EW} \sim 1$ TeV. Under some special assumptions, the model [6] leads to $\rho_V \sim (M_{EW}^2/m_{Pl})^4$. The numerical value of this combination of the energy scales is rather close, on the order of magnitude, to the observed figure [1,2].

In the present Letter, a framework is suggested that is alternative to the idea of quintessence. The cosmic acceleration is understood below as a dynamical effect of vacuum with constant density. Moreover, instead of time-dependent densities, constant time-independent numbers are introduced to the framework and identified as genuine physical characteristics of non-vacuum energies. The constants are among the basic quantities in cosmology; they are integrals of the Friedmann differential equations. For both non-relativistic and ultra-relativistic energies, the integrals express the conservation of the total number of particles in the comoving volume. It is interesting that the same equations attribute a special integral to vacuum as well, considering vacuum on a common basis with ordinary non-vacuum energies.

Two results are reported in the Letter. The first one comes from an empirical analysis of the problem. The evaluation of the cosmic integrals is made for vacuum, dark energy, baryons and ultra-relativistic energy with the use of the observed cosmic densities. It is found that the integrals are near-coincident numerically. Thus the cosmic coincidence appears in a novel version: this is now a coincidence of constant time-independent numbers (in contrast to the original version of the problem and its treatment in terms of quintessence). If this is not to be a numerical accident, the coincidence of the integrals looks like a new type of basic and stable regularities (a symmetry) in the evolving universe. Anyway, the integral coincidence exists as long as the energies themselves exist.

The second result comes from an attempt to clarify the physical nature of the integral coincidence on the grounds of fundamental physics. A simple kinetics theory model is developed for the freeze-out of non-relativistic relic particles in the early universe; this

is a well-know mechanism of the ‘origin of species’ in cosmology (see texts of Ref.[11]). The integrals enter the model naturally as its basic quantities. To specify the underlying physics, it is assumed that 1) the vacuum density is determined in terms of the electroweak energy scale and the Planck scale, as is proposed by the field theoretic model [6]; 2) the same two fundamental energy scales determine also the epoch of the freeze-out, so that the redshift at this epoch is $z \sim m_{Pl}/M_{EW}$; with this z , the temperature is $T \sim M_{EW} \sim 1$ TeV at the freeze-out. These assumptions are consistent with the central role played by electroweak physics in early cosmic evolution [11]. Under the two assumptions, the model demonstrates explicitly that the integrals for vacuum, non-relativistic and ultra-relativistic energies are coincident and have the correct numerical value. Thus the discovered coincidence of the cosmic integrals turns out to be a natural outcome of the freeze-out kinetics mediated by the electroweak scale physics at the cosmic age $\sim 10^{-12}$ sec.

Integrals. — The Friedmann cosmology equations,

$$a^{-2}(a')^2 = \kappa \rho a^2 - E, \quad d\rho = -3(\rho + p) d \ln a, \quad (1)$$

relate density and pressure to a , which is the curvature radius ($E = +1, -1$) or a scale factor ($E = 0$); the prime denotes a derivative with respect to conformal time, $\kappa = 8\pi G/3 = (8\pi/3)M_{Pl}^{-2}$, $E = -1, 0, +1$, in open, spatially flat and close models, respectively. With the Hubble constant $h_{100} = 0.65 \pm 0.15$, the observed densities lead to either an open model or a spatially flat model (see [3] and references therein).

For any cosmic energy having equation of state $p = w\rho$ with a constant parameter w , the integral of the second (‘thermodynamical’) equation in Eq.(1) is $C = \rho a^{3(1+w)} = Const > 0$. With this relation, the first (‘mechanical’) equation of Eq.(1) gives a as a function of time; in the resulting solution, integral C appears in a combination with the gravitational constant:

$$A = [(\frac{1+3w}{2})^2 \kappa C]^{\frac{1}{1+3w}} = [(\frac{1+3w}{2})^2 \kappa \rho a^{3(1+w)}]^{\frac{1}{1+3w}}. \quad (2)$$

In this form, integrals for different w have the same dimension (the length) and so can be compared with each other.

For the present epoch of accelerating expansion, the solution is

$$a(t) = A_V f(t), \quad f(t) = \sinh(t/A_V), \quad \exp(t/A_V), \quad \cosh(t/A_V), \quad (3)$$

for $E = -1, 0, +1$, respectively. Here $A_V = (\kappa\rho_V)^{-1/2}$, according to Eq.(2) with $w = -1$; the scale factor (for $E = 0$) is normalized to A_V . The acceleration is attributed to vacuum alone. The Hubble constant is $H \sim A_V^{-1}$, for each E soon after the transition to vacuum domination; the current cosmic age is $t_0 \sim H^{-1}$ in the transition era. As a result, one has approximately $a(t_0) \sim A_V$ for all the three models.

The integrals for dark (D) energy, baryons (B) and ultra-relativistic (R) energy with $w = 0, 0, 1/3$, respectively, enter corresponding cosmological solutions in the same quite natural way – see, for instance, an exact solution for both non-relativistic and ultra-relativistic energies [8].

The value of each of the four integrals can be found with the respective observed density and $a(t) = a(t_0)$; this way one has: $A_V = (\kappa\rho_V)^{-1/2} \sim 10^{61} M_{Pl}^{-1}$, $A_D = \frac{1}{4}\kappa\rho_D a^3 \sim 10^{60} M_{Pl}^{-1}$, $A_B = \frac{1}{4}\kappa\rho_B a^3 \sim 10^{59} M_{Pl}^{-1}$, $A_R = (\kappa\rho_R)^{1/2} a^2 \sim 10^{59} M_{Pl}^{-1}$. In the estimation of A_R , a conservative value $\alpha = 1$ is adopted which accounts for the CMB only. (Note that the integral for quintessence with $w = -2/3$, for example, is $A_Q \sim 10^{61} M_{Pl}^{-1}$, if it is estimated with $\rho_Q(t_0) = \rho_V$. A special case $w = -1/3$ is not included in Eq.(2); but using directly the corresponding solution of the Friedmann equations, one finds: $A_{-1/3} = A_V$.)

Thus the four cosmic integrals prove to be close to each other within two orders of magnitude:

$$A_w \sim 10^{60 \pm 1} M_{Pl}^{-1}, \quad w = [-1, 0, 0, 1/3]. \quad (4)$$

This is a novel version of the cosmic coincidence, which appears now as a tetrameric constant coincidence of constant numbers (in contrast to the temporary coincidence of the densities, in the original version of the problem). The coincidence of the two integrals A_B and A_R was recognized soon after the discovery of the CMB [9].

The integrals exist in the universe since the epoch at which the respective forms of energy start to exist themselves, and the integrals will exist until the decay of the protons at $t \sim 10^{31-32}$ sec and/or the decay of the particles of dark matter. Despite evolution

of the ratios of the densities over many orders of magnitude, the constant integrals are the same and coincident during all this extraordinarily large time interval.

As for the observed coincidence of the densities, it is clear that the densities of vacuum and dark matter are coincident, because this is the era of transition to the dynamical dominance of vacuum over dark energy, as is said above. The question of why we happen to live in such a special era may be treated on the ground of anthropic principle [10]. The coincidence of all the four major densities now is seen as a direct consequence of the basic relation of Eq.(4) and the temporary (and accidental, in this sense) coincidence $a(t_0) \sim A_A$ at the present epoch.

Model. — Is the observed coincidence of cosmic integrals a simple empirical fact or a reflection of a deeper physical regularity in nature? In a search for an answer to this question, one should address the physics of the early universe. To start with, one may consider D- and B-energies together as a non-relativistic thermal relic of early cosmic evolution. Then, one may consider the kinetics of freeze-out process in the early universe (see standard texts [11] and also a recent work [6]). As is well-known, for stable (or long-living) particles of the mass m , the abundance freezes out when the temperature T falls down the mass m , and the expansion rate $1/t$ wins over the annihilation rate, σn (here the annihilation cross-section $\sigma \sim m^{-2}$). At that particular moment, the particle density is $n \sim 1/(\sigma t) \sim m^2(G\rho_R)^{1/2}$. Using Eq.(2) for A_R and for $A_M = A_D + A_B$ with $\rho_M \sim mn$, one may now find how the two non-vacuum integrals are related in the model:

$$A_M \sim am^3 M_{Pl}^{-2} A_R. \quad (5)$$

One also has at that moment $\rho_R \sim m^4$, and because of this

$$A_R \sim a^2 m^2 M_{Pl}^{-1}. \quad (6)$$

Here and in Eq.(5) $a \simeq A_V(1+z)^{-1}$, and z is the redshift at the freeze-out epoch; in this way, the vacuum integral comes to the model.

To specify the underlying fundamental physics, one may refer again to the standard texts [11] that point out a special significance of electroweak physics in the early universe. If so, it is natural to identify the mass m of Eqs.(5,6) with the electroweak energy scale $M_{EW} \sim 1$ TeV. Then, one may exploit the field theoretic model [6] and adopt its

major result mentioned above, i.e. the relation for the vacuum density in terms of the two fundamental energy scales, M_{EW} and M_{Pl} : $\rho_V \sim (M_{EW}/M_{Pl})^8 m_{Pl}^4$. The only difference here from [6] is that the standard Planck scale is used, not the reduced Planck scale (see below). With this density, the vacuum integral is

$$A_V \sim (M_{Pl}/m)^4 M_{Pl}^{-1}. \quad (7)$$

Arguing along this line, one may expect that the epoch of the freeze-out is determined by the electroweak scale physics as well. If so, the redshift z at the freeze-out epoch may be a simple combination of the same two masses:

$$z \sim M_{Pl}/m. \quad (8)$$

Now the kinetics model is described by a system of four algebraic Eqs.(5-8) (with $m = M_{EW}$) for the four numbers A_M, A_R, A_V, z . The solution of the system is:

$$A_M \sim A_R \sim A_V \sim (M_{Pl}/m)^4 M_{Pl}^{-1}. \quad (9)$$

Thus the coincidence of the cosmic integrals appears as a natural result of the freeze-out process mediated by electroweak physics. This physics determines also the value of the integrals.

Following [6] and another recent work [12], one may introduce to Eq.(9) the gravitational scale M_G , or the reduced Planck scale m_{Pl} , instead of the standard Planck scale: $M_G \simeq m_{Pl} \simeq g M_{Pl}$, where $g \simeq 0.1 - 0.3$. The constant factor g accounts for the fact that the gravity constant G enters in combinations like $8\pi G/3, 6\pi G$, or $32\pi G/3$ exact formulas of cosmology. (Similarly, a few dimensionless factors, like the effective number of degrees of freedom, etc., may also be included in the model – see again Ref.[11]). So one gets finally:

$$A_w \sim g^4 (M_{Pl}/M_{EW})^4 M_{Pl}^{-1} \sim 10^{61 \pm 1} M_{Pl}^{-1}, \quad w = [-1, 0, 1/3]. \quad (10)$$

A quantitative agreement with the empirical result of Eq.(4) seems satisfactory here. With A_V of Eq.(10), one finds that the vacuum density is $\rho_A \sim g^8 (M_{Pl}/M_{EW})^8 M_{Pl}^4 \sim 10^{-122 \pm 2} M_{Pl}^4$, which is also in good agreement with the observed value [1,2] $\rho_A \sim 10^{-123} M_{Pl}^{-1}$.

Note that the numerical value of the redshift in Eq. (8) $z \sim gM_G/M_{EW} \sim 10^{15}$, and so the temperature at the freeze-out is $T \sim 1 \text{ GeV} \sim M_{EW}$, which is consistent with the central role of the electroweak scale in the model above.

Symmetry. — The tiny energy density of vacuum poses the most serious naturalness problem in theoretical physics [7]. With the identification of cosmic integrals as genuine physical quantities for both vacuum and non-vacuum energies, the naturalness problem finds its natural non-vacuum counterparts. In the framework suggested above, all the forms of cosmic energy can be treated in an unified manner and on the common ground of the cosmic integrals. So the naturalness problem is transformed, it loses its uniqueness and gets its place in a broader context that includes dark energy, radiation, etc. at the same rate. A more general problem is this: Why are $A_V \sim A_D \sim A_B \sim A_R \sim 10^{60} M_{Pl}^{-1}$?

Instead of the naturalness problem, one faces a new fundamental problem, which is the problem of symmetry that relates vacuum and non-vacuum forms of energy. The symmetry is described by Eqs.(4,10), in terms of cosmic integrals. This is translational symmetry in the one-dimensional space of energy forms, or of the values of the parameter w . Only three points of the space are directly observed at the phenomenological level: $w = -1, 0, 1/3$. The two more were mentioned above: $w = -2/3, -1/3$; each of them has a clear physical interpretation. The Zeldovich super-stiff equation of state, $w = 1$, may also be included to this collection. Quintessence is associated with a segment $-1 < w < -1/3$. The symmetry is not perfect, it turns out to be inexact, at a level of a few percent, on relative logarithmic scale.

Is the space of energy forms discrete or continuous? Is it really finite $-1 \leq w \leq 1$? Is the nature of the symmetry due completely to electroweak physics? Are quark-hadron physics, GUT, SUSY, etc. also mixed into making the symmetry inexact? These are the questions that are invited by the new appearance of the cosmic coincidence problem, and the list is obviously far incomplete.

The work was partly supported by the grant of the Academy of Finland.

- [1] S. Perlmutter *et al.*, *Astrophys. J* **517**, 565 (1998); A.G. Riess *et al.*, *Astron. J* **116**, 1009 (1998).
- [2] J. Cohn, astro-ph/9807128; S. Carol, astro-ph/0004075; N.Bahcall, J. Ostriker, S. Perlmutter, and P. Steinhard, *Science* **284**, 1481 (1999); O. Lahav, P.Lilje, J. Primack, and M. Rees, *MN RAS* **251**, 126 (1991); K.K.S. Wu, O. Lahav, and M.J. Rees, *Nature (London)* **397**, 225 (1999).
- [3] L. Wang, R.R. Caldwell, J.P. Ostriker, and P.J. Steinhardt, *Astrophis. J* **530**, 17 (2000).
- [4] R.R. Caldwell and P.J. Steinhardt, *Phys.Rev. D* **57**, 6057 (1998); B. Ratra and P.J.E. Peebles, *Phys. Rev. D* **37**, 3406 (1988) J.P. Ostraiker and P.J. Steinhardt, *Nature (London)* **377**, 600 (1995);
- [5] I. Zlatev, L. Wang, P.J Steihard, *Phys. Rev. Lett.* **82**, 896 (1999); A. Hebecker and C. Wetterich, *Phys. Rev. Lett.* **85**, 3339 (2000); V.A. Rubakov, *Phys. Rev. D* **61**, 061501 (2000); A.D. Dolgov, *Phys. Rev. D* **55**, 5881 (1997); C. Armandariz-Picon, V. Mukhanov, and P.J. Steinhard, *Phys. Rev. Lett.* **85**, 4438 (2000); I.G. Dymnikova, *Phys. Lett. B* **472**, 33 (2000); S. Dodelso nand M. Kaplinghat, E. Steward, *Phys. Rev. Lett.* **85**, 5276 (2000); A.Albrecht and C. Skordis, *Phys. Rev. Lett.* **84**, 2076 (2000).
- [6] N. Arkani-Hamed, L.J. Hall, Ch. Kolda, H. Murayama, *Phys. Rev. Lett.* **85**, 4434 (2000).
- [7] S. Weinberg, *Rev. Mod. Phys.* **61**, 1 (1989).
- [8] A.D. Chernin, *Astron. Zh.* **42**, 1124 (1965); Engl. transl. *Sov. Astron. -AJ* **9**, 871 (1966).
- [9] A.D. Chernin, *Pisma ZhETF* **8**, 633 (1968); Engl. transl. *Sov. Phys. - JETP Lett.* **8**, 391 (1968); A.D. Chernin, *Nature (London)* **220**, 250 (1968) - beware of typos.
- [10] S. Weinberg, *Phys. Rev. Lett.* **59**, 2607 (1987).
- [11] Ya.B. Zeldovich, I.D. Novikov, *The Structure and Evolution of the Universe*. The Univ. Chicago Press, Chicago and London (1983); A.D. Dolgov, Ya.B. Zeldovich, M.V. Sazhin, *Cosmology of the Early Universe (in Russian)*. Moscow Univ. Press, Moscow (1988); E.W. Kolb, M.S. Turner, *The Early Universe*. Addison-Wesley Pubs., New York (1990).

[12] M.Kawasaki, M. Yamaguchi, T. Yanagida, Phys. Rev. Lett. **85**, 3572 (2000).